

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/224873380>

Intergenerational change and familial aggregation of body mass index

Article in *European Journal of Epidemiology* · January 2011

CITATIONS

6

READS

18

9 authors, including:



[Jennifer Logue](#)

University of Glasgow

70 PUBLICATIONS 422 CITATIONS

[SEE PROFILE](#)



[Alex Mcconnachie](#)

University of Glasgow

264 PUBLICATIONS 4,924 CITATIONS

[SEE PROFILE](#)



[Niveen ME Abu-Rmeileh](#)

Birzeit University

89 PUBLICATIONS 2,341 CITATIONS

[SEE PROFILE](#)



[Naveed Sattar](#)

University of Glasgow

993 PUBLICATIONS 35,438 CITATIONS

[SEE PROFILE](#)

Johnson, P.C.D., Logue, J., McConnachie, A., Abu-Rmeileh, N.M.E., Hart, C., Upton, M.N., Lean, M.E.J., Sattar, N., and Watt, G.(2012) *Intergenerational change and familial aggregation of body mass index*. European Journal of Epidemiology, 27 (1). pp. 53-61. ISSN 0393-2990 (doi:10.1007/s10654-011-9639-5)

Copyright © 2012 Springer

A copy can be downloaded for personal non-commercial research or study, without prior permission or charge

Content must not be changed in any way or reproduced in any format or medium without the formal permission of the copyright holder(s)

When referring to this work, full bibliographic details must be given

<http://eprints.gla.ac.uk/58926/>

Deposited on: 05 November 2013

1 Title: Intergenerational change and familial aggregation of body mass index

2 Running title: Intergenerational change & familial aggregation of BMI

3 Authors:

4 Paul Johnson* PhD

5 Statistician

6 Robertson Centre for Biostatistics

7 Boyd Orr Building

8 University of Glasgow

9 Glasgow

10 G12 8QQ

11

12 Jennifer Logue* MBChB

13 Clinical Lecturer in Metabolic Medicine

14 Institute of Cardiovascular and Medical Sciences

15 University of Glasgow

16 126 University Place

17 Glasgow

18 G12 8TA

19

20 Alex McConnachie PhD

21 Senior Statistician

22 Robertson Centre for Biostatistics

23 Boyd Orr Building

24 University of Glasgow

25 Glasgow

26 G12 8QQ

27

28 Niveen M E Abu-Rmeileh PhD

29 Statistician

30 Institute of Community and Public Health

31 Birzeit University

32 Box 14

33 Birzeit

34 Occupied Palestinian territory

35

36 Carole Hart PhD

37 Reader in Public Health

38 Academic Unit of Public Health

39 Institute of Health and Wellbeing

40 University of Glasgow

41 1 Lilybank Gardens

42 Glasgow

43 G12 8RZ

44

45

46 Mark N Upton MBBS

47 General Practitioner

48 Woodlands Family Medical Centre

1 106 Yarm Lane
2 Stockton-on-Tees
3 Cleveland
4 TS18 1YE

5
6 Mike Lean MD
7 Professor of Human Nutrition
8 Institute of Health and Wellbeing
9 University of Glasgow
10 Glasgow
11 G12 8QQ

12
13 Naveed Sattar PhD
14 Professor of Metabolic Medicine
15 Institute of Cardiovascular and Medical Sciences
16 University of Glasgow
17 126 University Place
18 Glasgow
19 G12 8TA

20
21 Graham Watt MD
22 Professor of General Practice
23 Academic Unit of General Practice and Primary Care
24 Institute of Health and Wellbeing
25 University of Glasgow
26 1 Horselethill Road
27 Glasgow
28 G12 9LX

29
30 *Joint first authors

31
32 **Corresponding author**

33 Dr Jennifer Logue MBChB
34 Clinical Lecturer in Metabolic Medicine
35 BHF Glasgow Cardiovascular Research Centre
36 University of Glasgow
37 126 University Place
38 Glasgow
39 G12 8TA
40 Tel 0141 330 2569
41 Fax 0141 330 6955
42 Email Jennifer.Logue@glasgow.ac.uk

1 Funding: The Midspan Family Study was funded by the Wellcome Trust and the NHS
2 Cardiovascular Research and Development Programme. Neither the Wellcome Trust
3 nor the NHS Cardiovascular Research and Development Programme were involved in
4 the study design, the collection, analysis, and interpretation of data for this paper; in
5 the writing of the report; nor in the decision to submit the article for publication

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

1 **Abstract**

2 The relationship between parental BMI and that of their adult offspring, when
 3 increased adiposity can become a clinical issue, is unknown. We investigated the
 4 intergenerational change in body mass index (BMI) distribution, and examined the
 5 sex-specific relationship between parental and adult offspring BMI. Intergenerational
 6 change in the distribution of adjusted BMI in 1443 complete families (both parents
 7 and at least one offspring) with 2286 offspring (1263 daughters and 1023 sons) from
 8 the west of Scotland, UK, was investigated using quantile regression. Familial
 9 correlations were estimated from linear mixed effects regression models. The
 10 distribution of BMI showed little intergenerational change in the normal range (<25
 11 kg/m^2), decreasing overweightness ($25\text{--}<30 \text{ kg/m}^2$) and increasing obesity (≥ 30
 12 kg/m^2). Median BMI was static across generations in males and decreased in females
 13 by 0.4 (95% CI: $0.0, 0.7$) kg/m^2 ; the 95th percentile increased by 2.2 ($1.1, 3.2$) kg/m^2
 14 in males and 2.7 ($1.4, 3.9$) kg/m^2 in females. Mothers' BMI was more strongly
 15 associated with daughters' BMI than was fathers' (correlation coefficient (95% CI):
 16 mothers 0.31 ($0.27, 0.36$), fathers 0.19 ($0.14, 0.25$); $p=0.001$). Mothers' and fathers'
 17 BMI were equally correlated with sons' BMI (correlation coefficient: mothers 0.28
 18 ($0.22, 0.33$), fathers 0.27 ($0.22, 0.33$). The increase in BMI between generations was
 19 concentrated at the upper end of the distribution. This, alongside the strong parent-
 20 offspring correlation, suggests that the increase in BMI is disproportionately greater
 21 among offspring of heavier parents. Familial influences on BMI among middle-aged
 22 women appear significantly stronger from mothers than fathers.

23

24 **Keywords:** Obesity, body mass index, sex-specific, maternal, paternal.

25

1 **Introduction**

2 While there is no doubt that westernised populations are becoming more obese[1 2]
3 the distribution of weight change within the population is less well defined. The
4 majority of the published literature in this area uses data from the National Health and
5 Nutrition Examination Survey, based in the United States. Comparison of survey
6 results from 1994 with previous surveys in the late seventies and eighties showed that
7 the greatest absolute increases in body mass index (BMI) were within the already
8 heaviest group; significant upwards change in the entire distribution was only seen in
9 older age groups[3]. However, while such data shows trends in the BMI of the
10 population as a whole, studies describing changes within families are rare.

11
12 There has been increasing interest in the effects of parental BMI on childhood BMI,
13 encompassing the influence of genetics, maternal programming and environmental
14 factors; we have previously shown in this same cohort that midparental BMI is a
15 strong determinant of offspring BMI[4]. There is now evidence of a sex-specific
16 association, with one study reporting that childhood obesity is linked to obesity in the
17 same-sex parent[5], and another that both parents' BMI has an effect on offspring
18 BMI, but in female children the influence of mothers' BMI is stronger than
19 fathers'[6]. Those studies focused on an age when the child is still predominantly
20 dependent on their parents for nutrition, allowing potential for family-based
21 interventions. However the sex-specific link between parental obesity and obesity in
22 adult offspring, when any predisposition to obesity, including parity and sedentary
23 lifestyle, are likely to be expressed, and when increased adiposity more commonly
24 becomes a clinical issue, is unknown.

25

1 Using a two-generational study of 1443 sets of parents and 2286 adult offspring in the
 2 west of Scotland-based Midspan cohort, we examined for potential differences in
 3 BMI distribution between parents and offspring within the same population, using
 4 generational data gathered 20 years apart. The familial influences on BMI were
 5 examined by parent-offspring correlations, allowing the sex-specific relationship
 6 between parental and adult offspring BMI to be studied in depth.

7

8 **Methods**

9 *Study populations*

10 *The Midspan Renfrew/Paisley Study:* In 1972-76, 15402 residents of Renfrew and
 11 Paisley (7049 men and 8353 women), comprising 79% of the general population aged
 12 45-64 and including 4064 married couples, completed a questionnaire and attended
 13 for a clinical examination[7].

14

15 *The Midspan Family Study:* In 1993-4, current addresses were available for 3445
 16 couples from the Renfrew/Paisley Study (including the death certificate informant
 17 when both had died[8]); 2841 responded with information on the names, dates of birth
 18 and addresses of offspring. 3202 offspring from 1767 families were identified as
 19 living locally (within 30 miles), aged between 30 and 59 and therefore formed the
 20 eligible population for this study. In 1996 2338 offspring (1040 sons and 1298
 21 daughters) from 1477 families participated (73% response rate for individuals, 84%
 22 for families). In the present study excluding step-children, adopted offspring and
 23 families with a missing parental or offspring BMI, reduced the study sample to 1443
 24 complete families (both parents and at least one offspring included) with 2286
 25 offspring (1263 daughters and 1023 sons). The families were ascertained (by self

1 report) to be full-sibling families with no step-children, adoptees, half-sibs etc. All
2 were white. Details of the study have been described previously[4 8 9].

3

4 In addition to the 2338 participants in the Family study, there were 864 eligible
5 offspring who declined to take part. A further 1358 offspring were ineligible only
6 because they no longer lived locally. Sex, age, parental BMI and parental social class
7 were compared across these three groups (participants, local non-participants and
8 migrant non-participants) to investigate the possibility of migration and participation
9 bias.

10

11 *Physical measurements*

12 Standing height was measured in stockinged feet; in the offspring study a Holtain
13 stadiometer was used recorded to the nearest mm in 1996 and to the nearest cm in
14 1970s. Weight at both time points was measured to the nearest 0.1 kg in stockinged
15 feet and wearing indoor clothes. BMI was calculated as weight (kg)/ height (m^2), with
16 categories normal weight ($<25 \text{ kg/m}^2$), overweight ($25\text{-}29.9 \text{ kg/m}^2$) and obese (≥ 30
17 kg/m^2).

18

19 *Questionnaire*

20 Parents and offspring completed questionnaires recording marital status, smoking
21 status (never, current or former) and occupation. The offspring questionnaire also
22 asked for number of children. Marital status was recorded on the parent questionnaire
23 as married, single, widowed or other, and on the offspring questionnaire as married,
24 living with a partner, single, widowed, divorced or separated. Respondents who
25 identified themselves as married were classed as “married”; all others were classed as

“not married”. Social class was coded from occupation, using the Registrar General’s classification of occupation[10 11]. Social classes I, II, or III-nonmanual were defined as nonmanual, while III-manual, IV and V were defined as manual. Women’s social class was based on their own occupation or previous occupation, except housewives, where their husband’s or father’s occupation was used [9].

Statistical analyses

To allow comparison between parents and offspring, who differed in their distributions of age, marital status, number of children, smoking status and social class, all analyses were performed on BMI scores that had been adjusted to remove differences due to these potential confounding factors while preserving intergenerational differences. We used linear regression models to investigate the associations between BMI and potential confounding factors, separately for mothers, fathers, daughters and sons. The outcome was $\log(\text{BMI})$ and the explanatory variables were age (as a 3rd-order polynomial) (web figure 1), marital status, number of children (none, 1, 2, 3, ≥ 4), smoking status and social class (web figure 2). The residuals were added to the predicted $\log(\text{BMI})$ for a 50-year-old married never-smoker with two children and overall mean social class; taking exponentials gave adjusted BMI (BMI_{adj}).

BMI probability densities were estimated using a Gaussian kernel density estimator with bandwidth chosen by Silverman’s rule of thumb[12]. The mean, variance, and 5th, 25th, 50th, 75th and 95th percentiles of BMI_{adj} were estimated to assess intergenerational changes in the location and shape of the BMI distributions. Intergenerational change in percentiles of BMI_{adj} was estimated using quantile regression. Quantile regression models the relationship of the explanatory variables

1 with a given percentile of the outcome variable, in contrast with linear regression
 2 where the relationship with the mean of the outcome is modelled. Unlike linear
 3 regression, quantile regression allows us to investigate intergenerational change at
 4 specific points along the BMI_{adj} distribution. To illustrate the contribution of familial
 5 BMI to the BMI_{adj} distribution in the offspring generation, probability densities were
 6 estimated for offspring of normal weight and overweight parents.

7

8 Parent-offspring correlations in BMI were estimated from multilevel linear regression
 9 models. The standardised residuals from the models used to generate BMI_{adj}, as
 10 defined above, were denoted BMI-SDS (BMI standard deviation score). Multilevel
 11 models were fitted for BMI-SDS across all family members with separate family-level
 12 random intercepts for each family member group (pooled as parents and offspring or
 13 separated into mothers, fathers, daughters and sons). Within-family correlations were
 14 estimated as the correlation matrix of random intercepts.

15

16 The first model fitted assumed a common correlation between parents and offspring.
 17 Subsequent models assumed correlations to be sex-specific at the parental level (i.e.
 18 separate mother-offspring and father-offspring correlations), at the offspring level
 19 (separate parent-daughter and parent-son correlations) and at both levels (four
 20 separate correlation coefficients). When adjusting for potential confounders, missing
 21 social class for 68 subjects and missing number of children for two subjects were
 22 imputed using multiple imputation by additive regression. Statistical analyses were
 23 performed using the software packages MLwiN version 1.1[13] and R version
 24 2.10.0[14] with packages Hmisc and quantreg[15]. Multiple imputation was
 25 performed using the aregImpute R function[16].

1 **Results**

2 *Descriptive*

3 5172 subjects were included, comprising 1443 sets of parents, 1263 daughters and
 4 1023 sons. Table 1 allows intergenerational comparison between equivalent age
 5 cohorts by showing mean BMI and obesity prevalence divided into 5-year age bands.
 6 The mean (SD) parity was 2.9 (1.7) among the parents and 2.2 (0.9) among the 82%
 7 of offspring who were themselves parents. All the parents were married compared
 8 with 78% of the offspring. Parents were more likely to be current smokers (49% *vs*
 9 25%) and manual social class (61% *vs* 31%) compared with offspring. Web table 1
 10 describes the nonparticipants, both those living locally and those who migrated; there
 11 were no differences in parental BMI between participants and non-participants.

13 *Intergenerational change in BMI distribution*

14 The distribution of BMI_{adj} differed between generations in a number of ways (Table 2,
 15 Figure 1). Mean BMI_{adj} was 0.6 kg/m² (95% CI 0.3 to 0.9) higher in sons than in
 16 fathers, while mothers and daughters did not differ significantly. However median
 17 BMI_{adj} did not differ between generations in males, and decreased from mothers to
 18 daughters by 0.4 kg/m² (95% CI 0.0 to 0.7), suggesting that differences at the
 19 extremes of the distributions may be driving the difference between the means in
 20 males. Variance in BMI_{adj} increased across generations by 43% in females and 53% in
 21 males. In females the distribution of BMI_{adj} spread in both directions, less at the lower
 22 end but far more at the extreme upper end, while among males, the lower end and
 23 centre of the BMI_{adj} distribution were comparatively static across generations, while
 24 the upper end had increased. Since the age effect differed between generations in both
 25 sexes (Web Figure 1), the relatively small intergenerational changes detected in the

1 centre of the distribution (but not the larger increases observed in the upper tail) were
 2 sensitive to the choice of 50 years as the age to which to adjust BMI_{adj} (supplementary
 3 material).

4

5 *Familial influences on obesity and BMI*

6 The prevalence of obesity among offspring of normal weight parents (midparental
 7 BMI < 25 kg/m²) was 9%, compared with 24% among the children of overweight and
 8 obese parents (midparental BMI ≥ 25 kg/m²). Intra-familial correlations in BMI are
 9 reported in Table 3, derived from multilevel models of BMI-SDS. The parent-
 10 offspring correlation was 0.26 (95% CI: 0.23, 0.29). There was no difference between
 11 parent-daughter and parent-son correlations (p=0.423), suggesting that parental BMI
 12 predicts sons' and daughters' BMI equally well. There was strong evidence for a
 13 difference between mother-offspring and father-offspring correlations (p=0.016).
 14 However, this effect appears to be specific to daughters (p=0.001 for interaction): a
 15 mother's BMI is a better predictor than the father's BMI of their daughter's BMI.
 16 Maternal overweight and obesity is associated with a greater rightwards spread in the
 17 distribution of daughters' BMI than is paternal overweight and obesity (Figure 2).
 18 For sons, there is no evidence of a difference in correlation with mothers' and fathers'
 19 BMI (p=0.944): both parents' BMI is equally good at predicting their son's BMI
 20 (Web Figure 3).

21

22 We hypothesised that propensity for weight gain following childbirth might have
 23 contributed to the asymmetry between father-daughter and mother-daughter
 24 correlations. To test if parity had a role in weakening the father-daughter relative to
 25 the mother-daughter BMI correlation, we re-estimated the correlation coefficients in

1 Table 3 splitting daughters into those with (N=1065) and without children (N=198).
 2 The correlation (95% CI) between fathers and daughters with at least one child
 3 remained low at 0.18 (0.12, 0.24), while the correlation with childless daughters was
 4 0.23 (0.11, 0.36), slightly closer to the mother-daughter correlation. There was no
 5 significant difference between these correlations ($p=0.438$), and therefore no evidence
 6 for a role for parity, although the wide confidence interval for the difference (-0.08,
 7 0.19) suggests that this test has little power due to the low number of non-parous
 8 daughters.

9

10 Non-paternity would weaken the portion of father-offspring correlation that is due to
 11 shared genetic factors, and therefore could have contributed to the relatively weak
 12 father offspring correlations that we have observed. We investigated the impact of
 13 non-paternity by adjusting the correlations between mothers, fathers, daughters and
 14 sons under highly conservative assumptions of a 15% non-paternity rate and 100% of
 15 the father-offspring correlation being genetic[17 18]: the strength of the interaction
 16 was not substantially reduced (unadjusted $p=0.001$, adjusted $p=0.003$).

17

18 **Discussion**

19 This study adds to our understanding of the obesity epidemic in three ways. Firstly, it
 20 shows a pattern of changing adult body mass within one generation, characterised by
 21 the threshold defining the most overweight 5% of the population shifting substantially
 22 upwards (2-3 kg/m²), while the middle and lower portions of the distribution changed
 23 little (<1 kg/m²). Secondly, examination of familial influences on BMI showed that
 24 although both parents' BMI have an association with offspring adult BMI, maternal
 25 BMI is the significantly stronger influence on daughters' adult BMI, whereas both

1 parents influence sons' adult BMI equally. Finally, there is a very high prevalence of
2 obesity among adult offspring from overweight and obese parents compared with
3 offspring of normal weight parents (24% vs 9%).

4
5 The change in BMI distribution in this study confirms the findings of comparisons of
6 population based cross-sectional studies[19 20]: BMI has not increased evenly across
7 the population as a whole, but rather there has been a sharp increase in BMI at the
8 upper tail of the distribution. Broadly, the proportion of the population with normal
9 BMI is unchanged, while a decrease in overweightness is matched by a corresponding
10 increase in obesity. This pattern contradicts Rose's paradigm[21] of rising obesity
11 driven by a rightward shift in the entire distribution, but agrees with recent US cross-
12 sectional surveys[3] that suggest a "landslip" effect (Figure 1), where the overweight
13 are being replaced by the obese, but there is no corresponding recruitment into the
14 overweight cohort from those of normal weight.

15
16 The upward spread of the BMI distribution does not in itself imply that the increase is
17 concentrated among the most overweight families. Detection of such a trend is
18 complicated by regression to the mean, which predicts that the most overweight
19 parents will tend to have less overweight offspring[22]. However, purely artefactual
20 regression to the mean predicts stable variance across generations, while a genuine
21 tendency to divergence predicts increasing variance[23], as observed here. Thus the
22 upward spread of the BMI distribution over one generation, coupled with the positive
23 parent-offspring correlation, is consistent with the increase in BMI being
24 disproportionately among the adult offspring of heavier parents.

25

1 The sex-specific correlations observed here point to a substantial influence of shared
2 family environment on BMI, because no known genetic mechanism explains
3 daughters inheriting their BMI preferentially from their mothers. Relatively strong
4 mother-daughter BMI correlations (in this case relative to mother-son rather than
5 father-daughter correlations) were also found in a recent analysis of 4654 seven-year-
6 old children in the large ALSPAC cohort[6]. Another recent study of 226 children
7 aged 5-8 years in the EarlyBird cohort^[5] found both same-sex parent-offspring
8 correlations in BMI (i.e. both mother-daughter and father-son) but no significant
9 opposite sex correlations. We note that the EarlyBird analysis did not test for a
10 difference between same-sex and opposite-sex correlations, as was done here and in
11 the ALSPAC study, so the failure to find opposite sex correlations may have been a
12 consequence of small sample size rather than an indication of sex-specific inheritance.
13 Nevertheless, BMI category of same-sex parents was a better predictor of offspring
14 BMI than that of opposite sex parents[5]. In our analysis we have also adjusted for
15 potential confounders within both the parents and offspring, such as smoking, marital
16 and socioeconomic status, and number of children. We saw no difference for parental
17 effect on sons' BMI while daughters' were more strongly influenced by mothers'
18 BMI than fathers', and this difference was not explained by parity; however both
19 parents' BMI did have an effect on offspring BMI regardless of sex. Taken together,
20 these data are consistent with a model in which familial influences on daughters' BMI
21 are predominantly maternal in both young childhood and middle age, while familial
22 influences on sons' BMI is likely shared equally between both parents during
23 childhood and middle age.

1 It is widely accepted that parental BMI is related to offspring BMI[24]. Twin studies
2 have found BMI to be highly heritable[25 26] even in studies conducted during the
3 obesity epidemic[27], with a small environmental effect. However, genetics and
4 environment are closely linked in obesity; known obesity genes are thought to
5 increase susceptibility to obesity through control of food intake and food choice [28
6 29], hence why obesity has increased far faster than a genetic change would allow, as
7 the environment has changed. As women tend to do the majority of shopping and
8 cooking within a family, they have a strong influence over their children's diet; if they
9 are expressing their genotype by choosing high fat foods to feed the family, this may
10 explain why the mothers' influence is stronger. In this study however the offspring
11 were adults and there was no sex specificity over the sons' BMI; possible reasons for
12 this may include the influence of spouses on food provision and the influence of
13 fathers on sons' participation in sports and exercise. These results fit with the previous
14 findings in this cohort that parental socioeconomic position is more strongly
15 associated with offspring obesity in women than men[7 9 30]; there is a well
16 described socioeconomic gradient of environmental factors such as food choice and
17 availability that could be linked with increased obesity[31 32].

18 *Strengths and limitations*

20 The main strength of this study, in addition to the large sample size and the
21 availability of data on potential confounding factors, is the availability of adult
22 offspring. This unusual aspect of the study allowed familial influences on BMI to be
23 examined at the time when adiposity most often becomes clinically relevant. Another
24 strength is the positioning of the two generations on either side of a period of rapid
25 increase in obesity. However, environmental influences on obesity are likely to have

worsened further since 1996, so further research would be required to discover if the intergenerational patterns we have detected also apply to adults who are currently middle-aged. Other than that men were less likely to participate than females, no biases were found as a result of local offspring not participating in the study. There is a bias in the local eligible population towards the offspring of parents of manual social class and, possibly consequentially, mothers with higher BMI. If the trends in parental-offspring BMI correlations were similar in migrants and participants this BMI difference may have biased the results towards a larger right-shift in the offspring BMI; however, the size of the BMI difference between migrants and participants means the effect size would be very low.

Conclusions

Over one generation, the heaviest parents within our study population have been replaced by still heavier adult offspring while BMI in the remainder of the population has remained relatively unchanged. Strong parent-offspring correlations in BMI, even when the offspring are themselves adults, suggest that a large part of this increase has occurred within the heaviest families, possibly due to a combination of environmental, gestational and genetic influences. Further, we have shown for the first time that mothers appear to more strongly influence daughters' risk of obesity in adulthood than do fathers, indicating an environmental component alongside genetic factors.

COI statement: The authors declare no conflicts of interest.

Acknowledgements: Victor Hawthorne carried out the original Midspan studies.

Pauline MacKinnon is the Midspan administrator.

Ethics: Not required at the time of the Midspan Renfrew Paisley Study. Approval for the Midspan Family Study was granted from both the Argyll and Clyde, and Greater Glasgow Health Board Local Research Ethics Committees.

All authors contributed to the design of the study. PJ and AMcC analysed the data. PJ and JL wrote the first draft of the manuscript. All authors contributed to the redrafting of the manuscript and approved the final version. GW is the guarantor.

PJ and AMcC had full access to the data in the study and can take responsibility for the integrity of the data and the accuracy of the data analysis. The other authors had full access to all of the results.

Reference List

- 1 Government Office for Science. Foresight. Tackling obesities: future choices. 2010 Oct 17.
- 2 Ogden CL, Carroll MD, Curtin LR, et al. Prevalence of overweight and obesity in the United States, 1999-2004. *JAMA* 2006 Apr 5;**295**(13):1549-55.
- 3 Wang Y, Beydoun MA, Liang L, et al. Will all Americans become overweight or obese? estimating the progression and cost of the US obesity epidemic. *Obesity (Silver Spring)* 2008 Oct;**16**(10):2323-30.
- 4 Abu-Rmeileh NM, Hart CL, McConnachie A, et al. Contribution of Midparental BMI and other determinants of obesity in adult offspring. *Obesity (Silver Spring)* 2008 Jun;**16**(6):1388-93.
- 5 Perez-Pastor EM, Metcalf BS, Hosking J, et al. Assortative weight gain in mother-daughter and father-son pairs: an emerging source of childhood obesity. Longitudinal study of trios (EarlyBird 43). *Int J Obes (Lond)* 2009 Jul;**33**(7):727-35.
- 6 Leary S, Davey SG, Ness A. No evidence of large differences in mother-daughter and father-son body mass index concordance in a large UK birth cohort. *Int J Obes (Lond)* 2010 Jul;**34**(7):1191-2.
- 7 Hawthorne VM, Watt GC, Hart CL, et al. Cardiorespiratory disease in men and women in urban Scotland: baseline characteristics of the Renfrew/Paisley (midspan) study population. *Scott Med J* 1995 Aug;**40**(4):102-7.

- 1 8 Upton MN, McConnachie A, McSharpy C, et al. Intergenerational 20 year trends
2 in the prevalence of asthma and hay fever in adults: the Midspan family study
3 surveys of parents and offspring. *BMJ* 2000 Jul 8;**321**(7253):88-92.
- 4 9 Hart C, McConnachie A, Upton M, et al. Risk factors in the Midspan family
5 study by social class in childhood and adulthood. *Int J Epidemiol* 2008
6 Jun;**37**(3):604-14.
- 7 10 General Register Office. Classification of occupations 1966.London: HMSO,
8 1966. p. 1-148.
- 9 11 General Register Office. Classification of Occupations 1990.London: HMSO,
10 1990.
- 11 12 Silverman BW. *Density Estimation*. Chapman and Hall, 1986.
- 12 13 Rasbash J, Brown W, Healy M, et al. MLwiN Version 1.1. Multilevel Models
13 Project. 2011.
- 14 14 R Development Core Team. R: A language and environment for statistical
15 computing. 2009.
- 16 15 Roger Koenker. quantreg: Quantile Regression. R package version 4.50. 2010.
- 17 16 Harrell Jr FE. Hmisc: Harrell Miscellaneous. R package version 3.7-0. 2009.
- 18 17 Clemons T. A look at the inheritance of height using regression toward the
19 mean. *Hum Biol* 2000 Jun;**72**(3):447-54.
- 20 18 Davey SG, Steer C, Leary S, et al. Is there an intrauterine influence on obesity?
21 Evidence from parent child associations in the Avon Longitudinal Study of
22 Parents and Children (ALSPAC). *Arch Dis Child* 2007 Oct;**92**(10):876-80.
- 23 19 Flegal KM, Troiano RP. Changes in the distribution of body mass index of
24 adults and children in the US population. *Int J Obes Relat Metab Disord* 2000
25 Jul;**24**(7):807-18.
- 26 20 Micciolo R, Di F, V, Fantin F, et al. Prevalence of overweight and obesity in
27 Italy (2001-2008): is there a rising obesity epidemic? *Ann Epidemiol* 2010
28 Apr;**20**(4):258-64.
- 29 21 Rose G, Khaw KT, Marmot M. *Rose's Strategy of Preventative Medicine*.
30 Revised Edition ed. Oxford University Press, 2008.
- 31 22 Barnett AG, van der Pols JC, Dobson AJ. Regression to the mean: what it is and
32 how to deal with it. *Int J Epidemiol* 2005 Feb;**34**(1):215-20.
- 33 23 Hotelling H. Review of The triumph of mediocrity in business. *Journal of the*
34 *American Statistical Association* 1933;**28**:463-5.

- 24 Davey SG, Sterne JA, Fraser A, et al. The association between BMI and mortality using offspring BMI as an indicator of own BMI: large intergenerational mortality study. *BMJ* 2009;**339**:b5043.
- 25 Herskind AM, McGue M, Sorensen TI, et al. Sex and age specific assessment of genetic and environmental influences on body mass index in twins. *Int J Obes Relat Metab Disord* 1996 Feb;**20**(2):106-13.
- 26 Stunkard AJ, Harris JR, Pedersen NL, et al. The body-mass index of twins who have been reared apart. *N Engl J Med* 1990 May 24;**322**(21):1483-7.
- 27 Wardle J, Carnell S, Haworth CM, et al. Evidence for a strong genetic influence on childhood adiposity despite the force of the obesogenic environment. *Am J Clin Nutr* 2008 Feb;**87**(2):398-404.
- 28 Cecil JE, Tavendale R, Watt P, et al. An obesity-associated FTO gene variant and increased energy intake in children. *N Engl J Med* 2008 Dec 11;**359**(24):2558-66.
- 29 Bauer F, Elbers CC, Adan RA, et al. Obesity genes identified in genome-wide association studies are associated with adiposity measures and potentially with nutrient-specific food preference. *Am J Clin Nutr* 2009 Oct;**90**(4):951-9.
- 30 Power C, Graham H, Due P, et al. The contribution of childhood and adult socioeconomic position to adult obesity and smoking behaviour: an international comparison. *Int J Epidemiol* 2005 Apr;**34**(2):335-44.
- 31 Shelton NJ. What not to eat: inequalities in healthy eating behaviour, evidence from the 1998 Scottish Health Survey. *J Public Health (Oxf)* 2005 Mar;**27**(1):36-44.
- 32 Macdonald L, Cummins S, Macintyre S. Neighbourhood fast food environment and area deprivation-substitution or concentration? *Appetite* 2007 Jul;**49**(1):251-4.

1 Figure Legends

2

3 **Figure 1.** Estimated probability densities of adjusted BMI for parents (including those
4 with no same-sex offspring) and offspring, separately for males and females.

5

6 **Figure 2.** Probability density plots of adjusted BMI in daughters of normal weight
7 and overweight fathers and mothers. Percentiles are indicated by vertical lines.

8

9 **Web Figure 1.** Predicted average BMI given age (black line) \pm SE (grey line), in
10 mothers, fathers, daughters and sons. BMI was predicted from a model in which
11 log(BMI) was dependent on age (modelled as a 3rd-order polynomial), married status
12 (married, not married), number of children (none, 1, 2, 3, ≥ 4), smoking status (never,
13 former, current) and social class (manual, non-manual). Predictions were adjusted to a
14 married never-smoker with two children and mean social class (averaged across all
15 subjects). The age-BMI association is also represented by a LOESS smoothing line
16 (thin black line).

17

18 **Web Figure 2.** Predicted average BMI given married status, smoking habits, social
19 class and number of children. Predictions were adjusted as in Web Figure 1 for a 50-
20 year-old subject.

21 **Web Figure 3.** Probability density plots of adjusted BMI in sons of normal weight
22 and overweight fathers and mothers. Percentiles are indicated by vertical lines.

Table 1. Mean (SD) BMI and obesity prevalence by 5-year age bands in Midspan Family Study offspring and parents, and in 12,435 participants in the original Renfrew/Paisley Study who were not parents of Midspan Family Study offspring and for whom BMI was available.


		Age (years)								
		All	30-34	35-39	40-44	45-49	50-54	55-59	60-64	
Female	N	Daughters	1263	57	170	333	398	219	86	0
		Mothers	1443	0	0	0	436	516	314	177
		Renfrew/Paisley	6866	0	0	0	1585	1764	1723	1794
	Mean (SD) BMI (kg/m ²)	Daughters	25.9 (5.0)	25.5 (6.1)	25.6 (5.2)	25.6 (4.7)	26.2 (4.9)	25.7 (4.9)	26.7 (5.1)	-
		Mothers	25.9 (4.3)	-	-	-	25.4 (4.0)	25.5 (4.2)	26.8 (4.6)	26.8 (4.7)
		Renfrew/Paisley	25.7 (4.5)	-	-	-	25.2 (4.2)	25.4 (4.3)	25.7 (4.6)	26.5 (4.9)
	N (%) Obese (BMI ≥ 30 kg/m ²)	Daughters	229 (18.1%)	12 (21.1%)	33 (19.4%)	63 (18.9%)	68 (17.1%)	37 (16.9%)	16 (18.6%)	-
		Mothers	215 (14.9%)	-	-	-	58 (13.3%)	65 (12.6%)	66 (21.0%)	26 (14.7%)
		Renfrew/Paisley	1042 (15.2%)	-	-	-	188 (11.9%)	228 (12.9%)	265 (15.4%)	361 (20.1%)
Male	N	Sons	1023	56	157	263	317	168	62	0
		Fathers	1443	0	0	0	212	514	385	332
		Renfrew/Paisley	5571	0	0	0	1597	1452	1290	1232
	Mean (SD) BMI (kg/m ²)	Sons	26.5 (4.0)	25.8 (3.9)	25.8 (3.6)	26.5 (4.1)	27.0 (4.2)	26.8 (4.1)	25.9 (3.5)	-
		Fathers	26.0 (3.3)	-	-	-	25.8 (2.6)	26.2 (3.4)	26.0 (3.3)	25.8 (3.5)
		Renfrew/Paisley	25.8 (3.4)	-	-	-	25.9 (3.4)	25.9 (3.5)	25.8 (3.4)	25.8 (3.4)
	N (%) Obese (BMI ≥ 30 kg/m ²)	Sons	183 (17.9%)	9 (16.1%)	16 (10.2%)	48 (18.3%)	72 (22.7%)	29 (17.3%)	9 (14.5%)	-
		Fathers	161 (11.2%)	-	-	-	11 (5.2%)	69 (13.4%)	39 (10.1%)	42 (12.7%)
		Renfrew/Paisley	590 (10.6%)	-	-	-	168 (10.5%)	169 (11.6%)	130 (10.1%)	123 (10.0%)

Table 2. Intergenerational change in characteristics of the distributions of BMI_{adj} in females and males.

		Parent	Offspring	Difference (95% CI)	P-value
Female	N	1443	1263		
	Mean	26.3	26.4	0.2 (-0.2, 0.5)	0.380
	Variance	17.5	24.8	7.3 (5.0, 9.7)	<0.001
	5% percentile	20.5	20.0	-0.5 (-1.0, 0.0)	0.035
	25% percentile (Q1)	23.4	22.9	-0.4 (-0.8, -0.1)	0.021
	50% percentile (median)	25.8	25.5	-0.4 (-0.7, 0.0)	0.039
	75% percentile (Q3)	28.5	28.8	0.3 (-0.2, 0.9)	0.256
	95% percentile	33.5	36.1	2.7 (1.4, 3.9)	<0.001
Male	N	1443	1023		
	Mean	26.6	27.2	0.6 (0.3, 0.9)	<0.001
	Variance	10.7	16.3	5.6 (4.0, 7.2)	<0.001
	5% percentile	21.3	21.4	0.0 (-0.5, 0.6)	0.875
	25% percentile (Q1)	24.5	24.5	0.0 (-0.3, 0.3)	0.998
	50% percentile (median)	26.5	26.6	0.1 (-0.2, 0.5)	0.451
	75% percentile (Q3)	28.6	29.4	0.8 (0.4, 1.3)	0.001
	95% percentile	32.1	34.2	2.2 (1.1, 3.2)	<0.001

Table 3. Correlations (and 95% confidence intervals) between parent and offspring BMI, estimated from multilevel models where the response was BMI-SDS, and family relationships (mother, father, daughter and son) were fitted as random effects.

		Offspring			Daughter-son difference p-value
		Sons and daughters	Daughter	Son	
Parent	Mother and father	0.26 (0.23, 0.29)	0.25 (0.22, 0.29)	0.28 (0.24, 0.32)	0.423
	Mother	0.30 (0.26, 0.34)	0.31 (0.27, 0.36)	0.28 (0.22, 0.33)	
	Father	0.23 (0.19, 0.27)	0.19 (0.14, 0.25)	0.27 (0.22, 0.33)	
	Mother-father difference p-value	0.016	0.001	0.944	

**Figure 1**




Figure 2 (top)







Figure 2 (bottom)




Web Figure 1



Web Figure 2



Web Figure 3 (top)



Web Figure 3 (bottom)

Web Table 1. Characteristics of 4560 participant and non-participant offspring, with tests for migration bias (comparing local and migrant offspring) and response bias (comparing local participants and local non-participants). Differences in prevalence were tested using χ^2 tests and differences in means were tested using two sample t-tests.

		Local participant	Local non-participant	Migrant non-participant	N _{MISSING}	Migration bias P-value	Response bias P-value
N		2338	864	1358			
Sex	N (%) Male	1040 (44.5%)	475 (55.0%)	699 (51.5%)	0	0.010	<0.001
Age (years)	Mean (SD)	45.0 (6.2)	45.0 (6.8)	45.2 (6.2)	0	0.423	0.846
Maternal BMI (kg/m ²)	Mean (SD)	26.0 (4.4)	26.1 (4.5)	25.6 (4.3)	7	0.001	0.473
Paternal BMI (kg/m ²)	Mean (SD)	26.0 (3.3)	26.0 (3.5)	25.9 (3.2)	4	0.300	0.958
Maternal social class	N (%) Manual	1306 (57.8%)	493 (59.8%)	658 (50.5%)	174	<0.001	0.339
Paternal social class	N (%) Manual	1594 (68.9%)	606 (71.1%)	835 (61.9%)	43	<0.001	0.218

Supplementary Information

Sensitivity analysis for choice of age adjustment

Since the age effect differed between generations in both sexes (Web Figure 1), intergenerational comparisons were sensitive to the choice of 50 years as the age to which to adjust BMI_{adj}. Adjusting instead to 45 years had very little effect on intergenerational differences in male BMI_{adj} distribution, as might be expected by the fact that the age-BMI relationship in males is parallel across this age range (Web Figure 1). However, because there is a positive age effect in the mothers but none in the daughters, reducing the adjustment age to 45 years drew the mothers' distribution down to make the pattern of intergenerational change match closely to that of the fathers: no change in the left hand side of the distribution and a rightward stretch in the right hand side. Adjusting to 55 years had the opposite effect of shifting the BMI_{adj} distributions of both sons and daughters to the left, but not enough to counteract the stretching out of the upper tail. For all three adjustment ages the 95th percentile for BMI was significantly higher in offspring than in parents.